

A New Overgrowth Syndrome is due to Mutations in *RNF125*

Jair Tenorio,^{1,2} Alicia Mansilla,³ María Valencia,^{1,4} Víctor Martínez-Glez,^{1,5} Valeria Romanelli,¹ Pedro Arias,^{1,2} Nerea Castrejón,⁶ Fernando Poletta,⁷ Encarna Guillén-Navarro,^{1,8} Gema Gordo,² Elena Mansilla,^{1,9} Fé García-Santiago,^{1,9} Isabel González-Casado,¹⁰ Elena Vallespín,^{1,5} María Palomares,^{1,5} María A. Mori,^{1,5} Fernando Santos-Simarro,^{1,11} Sixto García-Miñaur,^{1,11} Luis Fernández,^{1,12} Rocío Mena,^{1,5} Sara Benito-Sanz,^{1,13} Ángela del Pozo,¹⁴ Juan Carlos Silla,¹⁴ Kristina Ibañez,¹⁴ Eduardo López-Granados,¹⁵ Alex Martín-Trujillo,¹⁶ David Montaner,^{1,17} The SOGRI Consortium,¹⁸ Karen E. Heath,^{1,13} Ángel Campos-Barros,^{1,13} Joaquín Dopazo,^{1,17} Julián Nevado,^{1,5} David Monk,¹⁶ Víctor L. Ruiz-Pérez,^{1,4} and Pablo Lapunzina^{1,2,11*}

¹CIBERER, Centro de Investigación Biomédica en Red de Enfermedades Raras, ISCIII, Madrid, Spain; ²Molecular Endocrinology Unit - Overgrowth Syndromes Laboratory, INGEMM, Instituto de Genética Médica y Molecular, IdiPAZ, Hospital Universitario La Paz, Universidad Autónoma de Madrid (UAM), Madrid, Spain; ³Instituto Cajal, CSIC, Madrid, Spain; ⁴Instituto de Investigaciones Biológicas, IIB, Universidad Autónoma de Madrid, Madrid, Spain; ⁵Functional and Structural Genomics Unit, INGEMM, Instituto de Genética Médica y Molecular, IdiPAZ, Hospital Universitario La Paz-UAM, Madrid, Spain; ⁶Developmental Cancer Group, Hospital San Juan de Dios, Barcelona, Spain; ⁷ECLAMC, Estudio Colaborativo Latinoamericano de Malformaciones Congénitas at CEMIC, Buenos Aires, Argentina; ⁸Medical Genetics Service, Hospital Virgen de la Arrixaca, Murcia, Spain; ⁹Cytogenetics Unit, INGEMM, IdiPAZ, Hospital Universitario La Paz-UAM, Madrid, Spain; ¹⁰Pediatric Endocrinology Unit, Hospital Universitario La Paz, Madrid, Spain; ¹¹Clinical Genetics Unit, INGEMM, IdiPAZ, Hospital Universitario La Paz-UAM, Madrid, Spain; ¹²Preanalytics Unit, INGEMM, IdiPAZ, Hospital Universitario La Paz-UAM, Madrid, Spain; ¹³Molecular Endocrinology Unit, INGEMM, IdiPAZ, Hospital Universitario La Paz-UAM, Madrid, Spain; ¹⁴Bioinformatics Unit, INGEMM, IdiPAZ, Hospital Universitario La Paz, Madrid, Spain; ¹⁵Department of Immunology, IdiPAZ, Hospital Universitario La Paz, Madrid, Spain; ¹⁶Laboratory of Epigenetics, Cancer Epigenetics and Biology Program, Idibell, Barcelona, Spain; ¹⁷Bioinformatics Unit, Centro de Investigación Príncipe Felipe, Valencia, Spain; ¹⁸SOGRI Consortium (Spanish OverGrowth Registry Initiative) Consortium at INGEMM, IdiPAZ, Hospital Universitario La Paz-UAM, Madrid, Spain

Communicated by Iain McIntosh

Received 4 June 2014; accepted revised manuscript 26 August 2014.

Published online 6 September 2014 in Wiley Online Library (www.wiley.com/humanmutation). DOI: 10.1002/humu.22689

ABSTRACT: Overgrowth syndromes (OGS) are a group of disorders in which all parameters of growth and physical development are above the mean for age and sex. We evaluated a series of 270 families from the Spanish Overgrowth Syndrome Registry with no known OGS. We identified one *de novo* deletion and three missense mutations in *RNF125* in six patients from four families with overgrowth, macrocephaly, intellectual disability, mild hydrocephaly, hypoglycemia, and inflammatory diseases resembling Sjögren syndrome. *RNF125* encodes an E3 ubiquitin ligase and is a novel gene of OGS. Our studies of the *RNF125* pathway point to upregulation of RIG-I-IPS1-MDA5 and/or disruption of the PI3K-AKT and interferon signaling pathways as the putative final effectors.

Hum Mutat 35:1436–1441, 2014. © 2014 Wiley Periodicals, Inc.

KEY WORDS: overgrowth; macrocephaly; autoimmune disorder; intellectual disability; *RNF125*

Generalized overgrowth syndromes (OGS) include the classic overgrowth disorders in which all of most parameters of growth and physical development are above the mean for age and sex [Lapunzina et al., 2002; Lapunzina, 2005]. Partial, localized, or regional OGS are those disorders in which excessive growth is confined to one or a few regions of the body. Macrocephaly is usually part of the clinical features observed in many OGS such as the Sotos (MIM #117550), Weaver (MIM #277590), Simpson–Golabi–Behmel (MIM #312870), Bannayan–Riley–Rubalcava (MIM #153480), Primrose (MIM #259050) [Cordeddu et al., 2014], and the 17q11.2 deletion (van Asperen syndrome; MIM #613675) [van Asperen et al., 1998] as well as in hemimegalencephaly-capillary malformation (MIM #602501) and CLAPO association (MIM #613089) [Lopez-Gutierrez and Lapunzina, 2008]. In 2007, Douglas et al. identified mutations in *RNF135* (an ubiquitin E3 ligase also known as Riplet mapping in the 17q11.2 deletion interval) in six families with a phenotype similar to the van Asperen syndrome but without neurofibromatosis [Douglas et al., 2007]. The authors demonstrated that haploinsufficiency of a RING finger gene was responsible for this OGS. All patients had macrocephaly (ranging from +2 to +4.1 SD), most were tall and heavy and some of them had relatives with mutations and similar clinical features. This clinical entity shares many clinical features with the van Asperen syndrome and has been recently renamed as macrocephaly–macroscopia–facial dysmorphism syndrome (MMFD; MIM #614192).

Additional Supporting Information may be found in the online version of this article.

*Correspondence to: Pablo Lapunzina, IdiPAZ - Instituto de Investigación Sanitaria del Hospital Universitario La Paz- Universidad Autónoma de Madrid-CIBERER- Centro de Investigación Biomédica en Red de Enfermedades Raras, ISCIII, Madrid, Spain; Paseo de la Castellana, 261-28046 Madrid, Spain. E-mail: pablo.lapunzina@salud.madrid.org

Contract grant sponsors: Fondo de Investigación Sanitaria (ISCIII [FIS 11/2491]); ENDOSCREEN (S2010/BMD-2396); The Marató-TV3; the EUCID-COST Project (BM1208); The Dravet Syndrome Foundation grant through the BBK-KutxaBank; The Spanish Sotos Syndrome Association.

Ubiquitin-proteasome degradation is one of the major post-transcriptional processes regulating the levels and function of proteins. The ubiquitin pathway includes the activity of at least three different enzymes: an ubiquitin-activating enzyme or E1, a conjugating enzyme or E2, and a ligase enzyme or E3. More than 600 E3 ubiquitin ligases are expressed in the human genome, allowing for the specificity of the ubiquitination system. However, the implication of the ubiquitin-proteasome pathway in OGS is not well known. E3 ubiquitin-ligases are involved, among others, in the phosphoinositide 3-kinase (PI3K) pathway. PI3K enzymes generate 3-phosphorylated phosphoinositides that act as second messengers downstream of numerous cellular receptors implicated in the regulation of cell growth, proliferation, survival, differentiation, and cytoskeletal changes. One of the best characterized targets of PI3K lipid products are the protein kinases AKT (v-AKT murine thymoma viral oncogene). The PI3K-AKT pathway has been implicated in the pathophysiology of several OGS such as the Proteus (MIM #176920), hemihyperplasia-hypoglycemia (240900), CLOVES (MIM #612918), and hemimegalencephaly-capillary malformation (MIM #602501) syndromes due to mutations in *AKT1*, *AKT2*, *AKT3*, *PI3KCA*, and/or *PI3KR2* [Lindhurst et al., 2011; Lindhurst et al., 2012; Riviere et al., 2012]. In this article, we report six patients from four families with overgrowth, macrocephaly and other findings, in whom we have found mutations in another RING finger gene, *RNF125* which encodes an E3 ligase. Similarly to RNF135, RNF125 is upstream of RIG-I (retinoic acid-inducible gene) and MDA5 (melanoma differentiation-associated gene 5), two proteins implicated in a variety of biological pathways including the interferons and the PI3K-AKT signaling pathways.

We selected a subset of 270 patients out of a total of 1,632 individuals registered at the Spanish Overgrowth Syndrome Registry with non-syndromic overgrowth (as defined by Neri in 2001) [Cohen et al., 2001]. We included all individuals with OGS who were negative for *NSD1*, *EZH2*, *GPC3*, and 11p methylation defects. We performed in all of them karyotypes and in most of them array CGH or SNP arrays (Supp. Materials and Methods). On SNP-array screening, one patient (patient 1, Fig. 1A and C, Supp. Table S1) showed a *de novo* 9 Mb deletion at chromosome 18 (chr18:24658770-34038769) encompassing candidate genes, including *RNF125*, *RNF138* and many other RefSeq genes (Supp. Table S2). This finding led us to redirect our investigations in the subset, by screening *RNF125* and *RNF138* in all 270 individuals using multiplex ligation probe amplification), high-resolution melting, Sanger sequencing, and pyrosequencing (Supp. Materials and Methods).

Three variants were identified in *RNF125* (c.336G>A [p.M112I], c.488C>T [p.S163L], and c.520C>T [p.R174C] (Fig. 1D and E) which were predicted, by in silico analysis, to be pathogenic (Supp. Table S3). In one of the patients, the mutation was *de novo* and in the other two patients the mutations were also observed in their affected mothers (patients 4 and 6, respectively) (Supp. Table S1, Fig. 1A and F). Pyrosequencing of *RNF125* in patients 2–6 confirmed that heterozygous point mutations were 1:1 in dosage and that there was no evidence of mosaicism (Fig. 1E). No pathogenic mutation in *RNF138* was found in any of the 270 patients. Clinical features and a summary of the studies performed in the four patients are listed in Supp. Table S1. The three missense mutations in *RNF125* were absent in 600 chromosomes from Spanish healthy controls and in 350 exomes of Spanish controls. We also explored the 1000 genome database (<http://browser.1000genomes.org>) and the Exome Variant Server (EVS) (<http://evs.gs.washington.edu/EVS/>). None of the three variants were reported in the 1000 genomes. In the EVS, we found that c.336G>A (p.M112I) was absent, whereas the other two variants were reported in 1/13005 chromosomes (0.0077%, Supp.

Table S3). Predictions using the PolyPhen2, Mut Pred, and Mutation Taster software suggest that all three changes are pathogenic or damaging (Supp. Table S3). Furthermore, methionine 112 and serine 163 residues are both highly conserved in mammals, whereas arginine 174 is conserved in almost all but a few mammals (Supp. Fig. S1). Protein modeling using Phyre2 predicted moderate to severe changes in the mutant proteins (Supp. Fig. S2).

To test other possible genetic or epigenetic alterations, we performed whole expression and whole methylation analysis (Supp. Figs. S3 and S4). We retrospectively also applied whole exome sequencing of DNA samples from blood of patients 2 and 5, to exclude the presence of other pathogenic overgrowth-causing mutations (Supp. Materials and Methods). After filtering for common variants, known polymorphisms and comparison with Spanish databases of exomes, bioinformatic analysis demonstrated that they shared variants in *RNF125* and seven other genes (Supp. Table S4). Segregation analysis in two affected mothers strongly suggested *RNF125* as the unique candidate gene (Fig. 1F). In all patients, we also applied whole-genome expression arrays and subsequently RT-qPCR confirmation in the three main pathways, RIG-I-IPS1, PI3K-AKT, and interferon signaling pathways, in which RNF125 participates. Clear differences were observed between patients and controls. The mRNA levels of *RNF125* were low in all patients (Fig. 2A) compared with controls, whereas *RNF135* levels were similar in the two groups (data not shown). In the RIG I-IPS1 pathway, some transcripts (*RIG-I*, *LGP2*, *MDA5*, and *IPS1*) were upregulated in patients compared to age and sex-matched controls (Fig. 2A). Other genes downstream of *RNF125* showed a non-homogeneous behavior: *DUBA*, *TRAF3*, and *TRAF6* were downregulated, *IRF3* and *TRADD* were upregulated and *TRAF2* was equal to controls (Supp. Fig. S4A). Similar results were obtained after the analysis of genes implicated in the PI3K-AKT pathway. Levels of *mTOR*, *PIK3R2*, *BTK*, *ILK*, *MYD88*, *PAK1*, *EIF4G1*, *HSPB1*, *PTEN*, and *TLR4* (Fig. 2A and Supp. Fig. S4B) were upregulated in patients compared with controls whilst *ATK2*, *AKT3*, *EIF4B*, *EIF4E*, *PIK3CA*, *PIK3R1*, and *RPS6KB1* were downregulated (Supp. Fig. S4B). Finally, analysis of 15 genes of the interferon pathway showed that only two genes, *IFNA8* and *IFNA14* were down-regulated (Supp. Fig. S4C), whereas the remaining 13 genes showed no differences in their expression profile pattern between patients and controls. These results confirm that mutations in *RNF125* lead to transcriptional downregulation of the gene and that haploinsufficiency of *RNF125* leads to upregulation of *RIG-I*, *IPS1*, and *MDA5* and misregulation of at least three main pathways (RIG-I-IPS1, PI3K-AKT, and Interferon). We tested whether expression of RIG-I protein is dysregulated in patients with *RNF125* mutations and found that basal RIG-I levels were not significantly different in B-lymphocytes cells of patients 1 and 2 compared with controls (data not shown). However, after treatment with poly (I:C), which acts as a double strand RNA-activator of the RIG-I signaling pathway, RIG-I induction levels were doubled in patients compared to controls (Fig. 2B and C). In order to demonstrate that RNF125 haploinsufficiency slows RIG-I degradation, we monitored the degradation kinetics of RIG-I after blocking translation with cycloheximide, a standard approach used to assay proteasome-mediated degradation. When protein synthesis was blocked, the turnover of endogenous RIG-I was slower in patient cells than in controls (Fig. 2D). After 12 hr post-cycloheximide treatment, patient-derived cells retained ~50% of the amount of RIG-I, whereas RIG-I was almost fully degraded in control cells (Fig. 2D–E).

Overgrowth, macrocephaly and intellectual disability (ID) are observed in a relatively small number of disorders [Lapunzina, 2005]. We here report six individuals from four families with mutations in *RNF125* and macrocephaly, overgrowth, enlarged ventricles, a




Figure 1. **A:** Facial phenotypes of patients 1, 2, 5, and 6. Correlation among the individuals and their position in the pedigree is in brackets. Note macrocephaly and large forehead in all patients and in some of them, a broad nose and synophrys. Patient 1 also presents a large mouth and a squared face, patient 2 has fleshy lips and upturned nose and patients 5 and 6 non-curved eyebrows and deep-set eyes. **B:** MRI of patient 2. Arrow indicates the anomaly of the Galenous' vein; note mild hydrocephalus and septum cavum pellucidum (white arrow). **C:** SNP-array of patient 1; the deleted region is shaded. The upper panel represents the log₂ ratio score (dosage) and the lower panel shows the haplotype plot showing loss of heterozygosity (LOH) in the deleted region. The deletion coordinates are in hg19. **D:** Comparison of the mutant and wild type chromatograms showing the *RNF125* mutations in the three probands (RefSeq NM_017831.3; data submitted to <http://www.lovd.nl/RNF125>). **E:** Pyrosequencing confirmation of each mutation showing a 1:1 ratio. **F:** Pedigrees of the four families showing cosegregation of the mutation with the phenotype in families 3 and 4. The pedigrees also include: genotype (+ = wt; M = mutant); y: (years), OFC: occipitofrontal circumference; H: height; and mutation detected for each case. NR: not recorded.

variable degree of intellectual disability and in some, hypoglycemia and recurrent inflammatory disease resembling Sjögren syndrome. Two patients inherited the mutations from their affected mothers and in the other two the mutations were *de novo*. The p.M112I and p.R174C mutations cosegregated with the phenotypes of their affected mothers. In silico analysis of the amino acid changes in *RNF125* indicates that they occurred in protein domains with important biological functions: p.M112I and p.R174C directly affect the first (C2HC) and third (C2H2) zinc finger domains of the protein, respectively (Supp. Fig. S1). The p.S163L mutation is located between the second and the third (C2H2) zinc finger domains, thus potentially affecting the intramolecular folding and the interactions among these zinc fingers [Giannini et al., 2008]. The introduction of a cysteine instead of an arginine at the third zinc finger disrupts the balance of the finger domain from C2H2 to C3H2 and potentially may also impair its function. In addition, in silico 3D analysis of the mutant proteins also suggest that the mutations may potentially alter the tertiary structure (Supp. Fig. S2), thus, affecting their functions. Finally, and very importantly, expression studies confirmed that all analyzed patients showed decreased *RNF125* RNA levels.

All our patients with *RNF125* mutations that resulted in downregulation of *RNF125* were shown to have upregulation of *RIG-I*, *MDA5*

and *IPS1* (Fig. 2A). *RNF125* functions as a negative regulator of *RIG-I* and *MDA5* signaling and a positive T cell regulator [Arimoto et al., 2007]. Dysregulation of *RIG-I* has been implicated in several infectious diseases and disorders of the immune system [Giannini et al., 2008]. Further, gain of function mutations of *IFIH1* (*MDA5*) have been recently described in patients with Aicardi-Goutieres syndrome (MIM #225750) featuring CNS anomalies and autoinflammatory diseases, as observed in our patients [Rice et al., 2014]. In accordance with our findings, Rice et al. (2014) proposed that mutations in *IFIH1* (*MDA5*) cause a gain of function of the protein leading to an increase in its binding with the RNA in the innate immune response. Similarly, haploinsufficiency of *RNF125* (which is upstream in the same pathway and regulates both *RIG-I* and *MDA5*) (Supp. Fig. S5) would lead to an increase of *MDA5* levels resulting in a gain of function of it and potentially dysregulating the immune response. In the interferon pathway, both *IFNA8* and *IFNA14* were also down-regulated. Transcriptional upregulation and increased protein levels of *RIG-I* and/or *IPS1* have been reported to play a role in the pathophysiology of autoinflammation [Britto et al., 2013]. Interestingly, some patients reported herein suffered from recurrent inflammation with a chronic pattern (Supp. Table S1). Patient 2 had several episodes of Raynaud phenomena, dry conjunctivitis and recurrent aphthous stomatitis and patient 4 suffered from continuous episodes




Figure 2. Functional analysis of RNF125. **A:** Expression arrays showing dysregulation of the RIG-I and PI3K-AKT pathways. Boxplot of mRNA expression of different genes included in the RIG-I and PI3K-AKT signaling pathways: *RNF125*; *RIG-I* (*DDX58*); *IPS-1*; *MTOR*; *PIK3R2*; *BTK*; *ILK*; and *LGP2*. Case = average of normalized mRNA levels of patients (P1: Patient 1; P2: Patient 2; P5: Patient 5) Control = average of normalized mRNA levels of controls. For each patient one age and sex matched control were used. The color range indicates the lowest (blue) to highest (red) expression. **B–E:** Western blot analysis of mutations in RNF125. RIG-I degradation is impaired in RNF125 haploinsufficiency derived cells. **B:** RIG-I levels were examined by Western blot of untreated cells from each patient (1 and 2) or the corresponding controls (C) and compared with cells treated with 100 μ g/ml poly (I:C) for 16 hr. **C:** Densitometry was undertaken for three independent experiments, as in Figure 2B. The percentage of RIG-I induction was plotted in a graph, after values were corrected for actin expression. Patient cells showed a major percentage of induction compared to control cells. **D:** Patients' cells (patients 1, 2, 5, and 6) were treated with poly (I:C) followed by addition of cycloheximide (CHX). Cells were harvested at the indicated times after addition of CHX and RIG-I levels analyzed by Western blot. **E:** Degradation kinetic plots of patients 1, 2, 5, and 6 versus controls showed a significant difference between the kinetics of the patients and controls.

of keratitis, conjunctivitis, limbitis, serous otitis, and pneumonia findings that are usually observed in the Sjögren syndrome (MIM #270150), a common autoimmune disease. We observed transcriptional upregulation of RIG-I and IPS1 at the RNA and protein level in our patients. Three E3 ubiquitin ligases, RNF125, RNF135, and TRIM25 target RIG-I for ubiquitination [Arimoto et al., 2007; Gack et al., 2007; Oshiumi et al., 2010] allowing RIG-I, and the three RIG I-like receptors, namely MDA5 [*IFIH1*], LGP2 (or DEXH box polypeptide 58 [*DHX58*], (MIM #608588) and IPS1 to interact in a very complex cascade of downstream activations (Supp. Fig. S5). IPS1 is activated by RIG-I and MDA5 which in turn both interacts with RNF125, RNF135 and TRIM25 [Oshiumi et al., 2010] (Supp. Fig. S5). The interaction between RIG-I and IPS1 is induced by TRIM25-mediated Lys63-linked polyubiquitination at Lys 172 of the RIG-I CARDs (caspase activation and recruitment domains) [Gack et al., 2007]. Previous studies demonstrated that RNF125 binds IPS1 as well as MDA-5 and was also ubiquitinated

[Arimoto et al., 2007] confirming that RNF125 has activity against RIG-I, MDA5, LGP2, and IPS1. These findings were also confirmed by the fact that the concentrations of MDA5 and IPS1 were reduced in cells overexpressing RNF125 ectopically and by a suppression of conjugation of ubiquitin to these proteins was observed by treatment with siRNA specific to RNF125 [Arimoto et al., 2007].

Shared features in our patients were overgrowth, macrocephaly with prominent and large forehead and some degree of ID; however, some other features were not observed in all individuals (Supp. Table S1). Our patients have phenotypic similarities with individuals with van Asperen syndrome (chromosome 17q11.2 deletion; MIM #613675) and MMFD (MIM #614192) due to mutations in *RNF135* [Douglas et al., 2007]. Although *RNF135* haploinsufficiency either to deletion or mutations leads to disease in humans, knockout of the mouse homolog, *rnf135*, does not cause any apparent developmental defects but rather severely reduced type I interferon, and abrogates

the activation of RIG-I and RIG-I C-terminal domain polyubiquitination [Oshiumi et al., 2010]. In addition, *rig-i* deficient mice are usually embryonic lethal, but interestingly, a few mice were born alive and died within 3 weeks after birth showing growth retardation (the reverse phenotype of overgrowth) [Kato et al., 2005]. Thus, mutations in *RNF125* and *RNF135* could lead to the same biological effect: haploinsufficiency of *RNF125* would prevent RIG-I binding thus, impairing IPS1 function while haploinsufficiency of *RNF135* would prevent activation of RIG-I and IPS1; similarly (Supp. Fig. S5). Despite not knowing the precise mechanism through which *RNF125* and *RNF135* haploinsufficiency results in the dysregulation of the RIG-I-IPS1 pathway, we have clearly demonstrated that the patients with *RNF125* mutations result in dysregulation of the two main activators of this pathway.

We have also observed transcriptional activation of some of the phosphatases of the PI3K pathway in patients with *RNF125* haploinsufficiency. There is considerable experimental evidence for the dysregulation of E3 ligases, PI3K signaling abnormalities, and altered growth. Abnormal E3 ligases and growth retardation is reported in Angelman (MIM #105830), Johanson–Blizzard (MIM #243800) and Kaufman oculocerebrofacial (MIM #615057) syndromes, among others. Additionally, some conditions with overgrowth have E3 ligases dysregulation. Firstly, synaptic overgrowth was observed in the absence of the E3 ubiquitin ligase “*highwire*” together with an increased PI3K signaling in motor neurons [Mozer and Sandstrom, 2012]. Secondly, a Ring Finger E3 responsible of Ecdysone receptor (a steroid hormone) degradation in *Drosophila melanogaster* acts at high concentrations as a developmental timer and at basal levels contributes to the inhibition of tissue growth [Gradilla et al., 2011]. Thirdly, another E3 ligase complex associated to akt (*SCF^{slimb}*) inhibits neuroblast overgrowth in *drosophila* brains [Li et al., 2014]. Fourthly, the Big Brother (*bb*) gene, encoding an E3 ubiquitin ligase, represents a central regulator of organ size and growth in *Arabidopsis*. Plants lacking *bb* activity form larger organs; whereas conversely, plants expressing higher levels of *bb* produce smaller organs, indicating that *bb* acts as a negative regulator for organ size [Breuninger and Lenhard, 2012]. *RNF125* has a 46% (20/45) positive identity at the zinc finger domains with *bb* (amino acids 197–241). Exact amino acid identity between *RNF125* and *BB* occurs at the eight conserved cysteines and histidines of the zinc finger domains. Fifthly, individuals with haploinsufficiency of *NSD1* (a gene whose protein has several zinc fingers) develop Sotos syndrome (MIM #117550), an overgrowth/macrocephaly disorder. Thus, haploinsufficiency of E3 ubiquitin ligases in plants (*Arabidopsis*) and *D. melanogaster* leads to overgrowth and in humans, haploinsufficiency of *RNF135* [Douglas et al., 2007] and now in this study, *RNF125*, leads to overgrowth, macrocephaly and intellectual disability. Finally, firm evidence of the implication of the PI3K-AKT pathways in overgrowth-macrocephaly syndromes and links among the PI3K and the RIG-I/interferon pathway have recently been communicated. Mutations in *PIK3CA*, *PIK3R2* and *AKT3* are observed in the megalencephaly-capillary malformation [Riviere et al., 2012], in *AKT1* in Proteus syndrome [Lindhurst et al., 2011] and in *AKT2* in hemihypertrophy–hypoglycemia [Lindhurst et al., 2012]. In addition, PI3K has been defined as one of the main factors activated through RIG-I mediated anti-pathogen response to enhance expression of type I interferons. Other features observed in some of our patients were intellectual disability, mild hydrocephaly, and hypoglycemia. Intellectual disability and hydrocephaly are also observed in patients with *AKT3*, *PI3KR2*, and *PI3KCA* mutations, whereas hypoglycemia has been observed in individuals with *AKT2* mutations [Lindhurst et al., 2011; Lindhurst et al., 2012; Riviere et al., 2012]. Intellectual disability has been also described in patients with

mutations in other RING finger genes such as *RNF133* and *RNF148* [Bonora et al., 2012].

In summary, we have identified six affected individuals from four OGS families with mutations in *RNF125*. We have demonstrated that mutations in *RNF125* result in a loss of function of *RNF125* and dysregulation of the RIG-I-IPS1, PI3K-AKT, and interferon pathways. The exact mechanism/s by which haploinsufficiency of *RNF125* results in overgrowth, macrocephaly and in some patients intellectual disability, hypoglycemia, and Sjögren-like syndrome remains elusive. Though the complex pathways linking E3 ubiquitin ligases, RING fingers, CARD/CARD-like proteins and the RIG I-IPS1, PI3K-AKT and interferon pathways need further investigations, this article demonstrates the connections among these proteins and their implication in a novel disorder of growth, development and autoimmunity.

Acknowledgments

The SOGRI (Spanish OverGrowth Registry Initiative) Consortium is constituted by the following researchers: A. Plasencia, Alberto L. Rosa, Aleixandre Blanquer, Alfredo García-Alix, Alfredo Santana, Alicia Delicado, Almudena Alonso, Amaya Rodríguez, Amparo Sanchis, Ana Moreno, Ana Patiño García, Ana Vega, Analía Bredani, Andrea Paula Solari, Andrea Villavicencio, Angelina Acosta, Anibal Nieto, Anna Maria Cueto González, Antonio Baldellón, Antonio González Meneses, Antonio Martínez Carrascal, Aranzazu Díaz de Bustamante, Arteché Ocasar, Blanca Gener, Blasco González, Boris Groisman, Bradford Coffee, Carlos Alcalde Martín, Carmen Aragon Fernandez, Carmen Benito, Carmen Martín Seisdedos, Carmen Roche, Claudia Arberas, Claudia Perandones, Claudio Contessotto, Cristina Olivas, Daniel Armenta, Denise Cavalcanti, Dolores Elorza, Eduardo Castilla, Elena Zamora, Elisa Zambrano, Elisabeth Steichen, Enrique Caro Cruz, Enrique Galán Gómez, Enriqueta Román, Ernesto Goldschmidt, Esteban Marfil, Esther Gean, Eugenia Antolín, F. Javier Gascón Jiménez, Feliciano Ramos, Fermina López Grondona, Fernández Córdoba, Fernando Regla Vargas, Francisco Martínez, J. Miguel García Vegada, Giovannucci Uzielli, Gloria Gacio, Carmen González Armengod, Graciela Mercado, Hamilton Cassinelli, Ieda Orioli, Ignacio Arroyo, Ignacio Díez López, Ignacio Onsurbe Ramírez, Ignacio Pascual Castroviejo, Ignacio Pascual Pascual, Ignacio Vázquez Río, Inés Bueno, Isabel Espejo Portero, Isabel Lorda Sánchez, Jaime Sánchez del Pozo, Jaume Campistol, Javier Arcas, Javier Fernández, Javier García Planells, Javier López Pisón, Jesús Barreiro, Jesús del Valle Núñez, María José Jiménez Liria, Joaquín Fernández Toral, Joaquín Ramírez, Jordi Rosell, Jorge Vilaplana, José Carlos Cabral de Almeida, José Ignacio Labarta, José L. Herranz, José Luis Fernández Luna, José Luis Fuster, José M. Díaz, Jose M. Gairi, José Miguel García Sagredo, Juan A. Piñero, Juan Carlos López Gutiérrez, Juan Manuel Fernández, Juan P. López Siguero, Juan Tovar, Judith Armstrong, Julián Lara, Leonor Aranz, Laura Rodríguez, Leandro Soriano, Leila Cardoso, Liliana De Alba, Loreta Cimalistiene, Loreto Martorell, Luis González Gutiérrez Solana, Luis Pérez Jurado, M Asunción López Ariztegui, M. Antonia Molina, M. Cruz García, M. Ferrer Lozano, M. Jesús Alija Merillas, M. Luisa Martínez-Frías, María L. Martínez Fernández, M. Rocío Jadraque, M^a Asunción García Pérez, M^a Montserrat Rodríguez Pedreira, M^a Pilar Ribate, M^a Teresa González López, M^a Teresa Moral Pumarega, Mabel Segovia, Macarena Lizama, Manuel Pombo, Margarita Martínez, Margarita Taberner, María Antonia Ramos, María Ballesta, María Belar, María Jesús Lautre, Marta Cruz, M. Nieves Martínez Guardia, F. Javier Martínez Sarries, Mercedes Artigas, Mercedes Villanueva, Meritxell Torrabias, Miguel del Campo, Miguel Tomás Vila, Miguel Urioste, Mónica Rosello, Nik Kantaputra, Pablo Prieto Matos, Paloma Dorao, Paula Casano, Paula Lalaguna Mallada, Pedro Olivares, Raquel Perez Delgado, Priscila Bernardi, Rafael Camino León, Ramón Cañete, Ramón Gaztañaga, Ramón Velazquez, Ramón Vidal Samahuja, Raquel Sáez Villaverde, Ricardo Gracia, Richard Scott, Rita Valdez, Rosa Arteaga, Rosa Cedeño, Rosario Cazorla, Rosario Marín Iglesias, Rubén Bronberg, Salvador Climent, Santiago Conde Barreiro, Seema Kapoor, Soledad Kleppe, Sonia Santillán, Trinidad García Lopez, Teresa Calvo, Teresa Vendrell, Pilar Tirado, Claudia Toledo Pacheco, Alicia Ureta Huertos, Vanesa Lopez, Vanesa Lotersztejn, Vanesa Méndez,

Selma Vázquez Martín, Verónica Seidel, Vicente Albiach, Víctor M. Navas López, Virginia Soler, and Viviana Cosentino. We also thank the CIBERER BioBank for the support and all participating families.

References

- Arimoto K, Takahashi H, Hishiki T, Konishi H, Fujita T, Shimotohno K. 2007. Negative regulation of the RIG-I signaling by the ubiquitin ligase RNF125. *Proc Natl Acad Sci USA* 104:7500–7505.
- Bonora E, Wischmeijer A, Minopoli F, Tabarroni C, Vidone M, Seri M, Graziano C, Romeo G. 2012. Analysis of ring finger proteins RNF133 and RNF148 in intellectual disability. *Eur J Hum Genet* 20:Suppl. 1–78.
- Breuninger H, Lenhard M. 2012. Expression of the central growth regulator BIG BROTHER is regulated by multiple cis-elements. *BMC Plant Biol* 12:41.
- Britto AM, Amoedo ND, Pezzuto P, Afonso AO, Martinez AM, Silveira J, Sion FS, Machado ES, Soares MA, Giannini AL. 2013. Expression levels of the innate response gene RIG-I and its regulators RNF125 and TRIM25 in HIV-1-infected adult and pediatric individuals. *AIDS* 27:1879–1885.
- Cohen MM, Neri G, Weksberg R. 2001. *Overgrowth syndromes*: Oxford University Press, New York, NY, USA.
- Cordeddu V, Redeker B, Stellacci E, Jongejan A, Fragale A, Bradley TE, Anselmi M, Ciolfi A, Cecchetti S, Muto V, Bernardini L, Azage M, et al. 2014. Mutations in ZBTB20 cause Primrose syndrome. *Nat Genet* 46:815–817.
- Douglas J, Cilliers D, Coleman K, Tatton-Brown K, Barker K, Bernhard B, Burn J, Huson S, Josifova D, Lacombe D, Malik M, Mansour S, et al. 2007. Mutations in RNF135, a gene within the NF1 microdeletion region, cause phenotypic abnormalities including overgrowth. *Nat Genet* 39:963–965.
- Gack MU, Shin YC, Joo CH, Urano T, Liang C, Sun L, Takeuchi O, Akira S, Chen Z, Inoue S, and Jung JU et al. 2007. TRIM25 RING-finger E3 ubiquitin ligase is essential for RIG-I-mediated antiviral activity. *Nature* 446:916–920.
- Giannini AL, Gao Y, Bijlmakers MJ. 2008. T-cell regulator RNF125/TRAC-1 belongs to a novel family of ubiquitin ligases with zinc fingers and a ubiquitin-binding domain. *Biochem J* 410:101–111.
- Gradilla AC, Mansilla A, Ferrus A. 2011. Isoform-specific regulation of a steroid hormone nuclear receptor by an E3 ubiquitin ligase in *Drosophila melanogaster*. *Genetics* 189:871–883.
- Kato H, Sato S, Yoneyama M, Yamamoto M, Uematsu S, Matsui K, Tsujimura T, Takeda K, Fujita T, Takeuchi O, and Akira S. et al. 2005. Cell type-specific involvement of RIG-I in antiviral response. *Immunity* 23:19–28.
- Lapunzina P. 2005. Risk of tumorigenesis in overgrowth syndromes: a comprehensive review. *Am J Med Genet C Semin Med Genet* 137C:53–71.
- Lapunzina P, Camelo JS, Rittler M, Castilla EE. 2002. Risks of congenital anomalies in large for gestational age infants. *J Pediatr* 140:200–204.
- Li S, Wang C, Sandanaraj E, Aw SS, Koe CT, Wong JJ, Yu F, Ang BT, Tang C, Wang H. 2014. The SCFSlimb E3 ligase complex regulates asymmetric division to inhibit neuroblast overgrowth. *EMBO Rep* 15:165–174.
- Lindhurst MJ, Parker VE, Payne F, Sapp JC, Rudge S, Harris J, Witkowski AM, Zhang Q, Groeneveld MP, Scott CE, Daly A, Huson SM, et al. 2012. Mosaic overgrowth with fibroadipose hyperplasia is caused by somatic activating mutations in PIK3CA. *Nat Genet* 44:928–933.
- Lindhurst MJ, Sapp JC, Teer JK, Johnston JJ, Finn EM, Peters K, Turner J, Cannons JL, Bick D, Blakemore L, Blumhorst C, Brockmann K, et al. 2011. A mosaic activating mutation in AKT1 associated with the Proteus syndrome. *N Engl J Med* 365:611–619.
- Lopez-Gutierrez JC, Lapunzina P. 2008. Capillary malformation of the lower lip, lymphatic malformation of the face and neck, asymmetry and partial/generalized overgrowth (CLAPO): report of six cases of a new syndrome/association. *Am J Med Genet A* 146A:2583–2588.
- Mozer BA, Sandstrom DJ. 2012. *Drosophila* neuroigin 1 regulates synaptic growth and function in response to activity and phosphoinositide-3-kinase. *Mol Cell Neurosci* 51:89–100.
- Oshiumi H, Miyashita M, Inoue N, Okabe M, Matsumoto M, Seya T. 2010. The ubiquitin ligase Riplet is essential for RIG-I-dependent innate immune responses to RNA virus infection. *Cell Host Microbe* 8:496–509.
- Rice GI, del Toro Duany Y, Jenkinson EM, Forte GM, Anderson BH, Ariaudo G, Bader-Meunier B, Baildam EM, Battini R, Beresford MW, Casarano M, Chouchane M, et al. 2014. Gain-of-function mutations in IFIH1 cause a spectrum of human disease phenotypes associated with upregulated type I interferon signaling. *Nat Genet* 46:503–509.
- Riviere JB, Mirzaa GM, O’Roak BJ, Beddaoui M, Alcantara D, Conway RL, St-Onge J, Schwartzentruber JA, Gripp KW, Nikkel SM, Worthylake T, Sullivan CT, et al. 2012. De novo germline and postzygotic mutations in AKT3, PIK3R2 and PIK3CA cause a spectrum of related megalencephaly syndromes. *Nat Genet* 44:934–940.
- van Asperen CJ, Overweg-Plandsoen WC, Cnossen MH, van Tijn DA, Hennekam RC. 1998. Familial neurofibromatosis type 1 associated with an overgrowth syndrome resembling Weaver syndrome. *J Med Genet* 35:323–327.